

Dr Caroline Dowling

Urological Surgeon MS, FRACS (Urol)



Malvern Hill Consulting
1039 Malvern Road,
Toorak VIC 3142

Phone: 03 9188 3690
Fax: 03 9824 6991
Call Service 03 9387 1000
Web: www.urologyworks.com.au

Pubovaginal Slings

There are many surgical options for women with stress urinary incontinence (exertion related leakage, i.e. if you exert yourself, you leak) who have not responded to physiotherapy or medications. Pubovaginal slings are offered to women with stress incontinence who need support to the urethra (tube through which the urine passes from the bladder to the outside) to stay dry. This operation can be performed alone or in combination with other procedures such as vaginal repair or hysterectomy.

A fascial pubovaginal sling is usually recommended when the incontinence is severe or when the patient is young or we *wish to avoid using a synthetic material such as is used in the mid urethral slings*. There are some conditions also where the fascial sling is used to protect another repair that has been performed as in where there is an abnormality of the urethra.

- It is very effective, i.e. >90% cure rate for stopping stress incontinence
- There are very few associated problems
- After four weeks patients can return to normal activity, including sport of any kind
- Hospital stay required - 2-4 days
- A general anaesthetic is generally given. Occasionally a regional anaesthetic is used but the choice of anaesthesia is entirely at the discretion of the anaesthetist

How do I know if this operation is for me?

Urodynamic testing is done to confirm the diagnosis and that there is no other cause for your incontinence. This test will also help your surgeon make an informed decision with you about the suitability of this operation for you.

What Occurs During The Operation

This operation involves a vaginal and abdominal (on your tummy, low down just above the pubic hair) incision. A strip of tough material is taken from the covering of your abdominal muscles. This is called *rectus fascia*. That fascia is then tied to a special string type of material to enable the sling to be brought around the urethra in such a way that, when you cough and move, the sling tightens and closes the urethra to stop leakage.

What To Expect Post Operatively

- An abdominal wound is required – usually 8 cm – therefore lifting and activity involving the abdominal muscles needs to be limited for at least four weeks after surgery.
- In the first two weeks post operatively, difficulty emptying the bladder is common although some women pass urine without difficulty. Because of this

Dr Caroline Dowling

Urological Surgeon MS, FRACS (Urol)



Malvern Hill Consulting
1039 Malvern Road,
Toorak VIC 3142

Phone: 03 9188 3690
Fax: 03 9824 6991
Call Service 03 9387 1000
Web: www.urologyworks.com.au

recognised short term problem, all patients are taught how to pass a little catheter. This enables them to be free of the urinary catheter. Long term voiding trouble is rare but can occur. In less than 5% of patients this would need a loosening or cutting of the sling.

- Sling related discomfort, i.e. discomfort relieved by bending the knees up, is not uncommon in the short term after the sling. This usually resolves gradually over the weeks following surgery and no other sling discomfort is associated.
- The operation cures exertion related leakage. It does not cure all associated urge (the strong desire to pass urine) incontinence and sometimes, especially if the incontinence was severe, urge incontinence may occur for the first time. This usually settles and is treatable.
- There is usually some vaginal discharge for 2-3 weeks while the stitches are dissolving. After that intercourse is possible again depending on how comfortable you otherwise feel.

Can there be any complications from the procedure?

- A small risk of entering the bladder, urethra or blood vessels when the sling is passed from the abdomen to the vagina. Rarely this would require further surgery for this problem or any other reason encountered at the time of sling placement.
- Between one and five woman in a hundred will have trouble going to the toilet and may need a catheter for a short period after the surgery until normal bladder emptying is established
- Uncommonly division or adjustment of the tape is required (approx. 1 in 70).
- Between five and ten women in a hundred will develop an irritable bladder which usually improves after 1-3 months. Occasionally urgency and urge incontinence may be worse requiring medication
- Urine infection requiring antibiotic treatment.

Recovery time

Most patients return home after 2-4 days, once you are feeling well and are passing urine with no problem. If you need pain relief, tablets are usually enough. It is important to rest after the operation and allow the area to heal. Generally it is recommended:

- You restrict activity for two weeks
- After two weeks do light activity only
- Avoid heavy lifting, including shopping bags, washing baskets and children, for six weeks and ideally limit this to <10kg for three months
- Abstain from sexual activity for 4-6 weeks
- Avoid playing sport for four weeks.
- NO driving for 2-4 weeks